

Publicly Funded Immunization Schedules for Ontario – December 2016

Publicly funded vaccines may be provided only to eligible individuals and must be free of charge

Routine Schedule: Children Starting Immunization in Infancy													
Vaccine	Age	2 Months	4 Months	6 Months	12 Months	15 Months	18 Months	4-6 Years^	Grade 7	14-16 Years§	24-26 Years †	≥34 Years ‡	65 Years
DTaP-IPV-Hib Diphtheria, Tetanus, Pertussis, Polio, <i>Haemophilus influenzae</i> type b		◆	◆	◆			◆						
Pneu-C-13 Pneumococcal Conjugate 13		◆	◆		◆								
Rot-1 Rotavirus		▲	▲										
Men-C-C Meningococcal Conjugate C					◆								
MMR Measles, Mumps, Rubella					■								
Var Varicella						■							
MMRV Measles, Mumps, Rubella, Varicella								■					
Tdap-IPV Tetanus, diphtheria, pertussis, Polio								◆					
HB Hepatitis B									●				
Men-C-ACYW Meningococcal Conjugate ACYW-135									●				
HPV-4 Human Papillomavirus									●				
Tdap Tetanus, diphtheria, pertussis										◆	◆		
Td (booster) Tetanus, diphtheria												◆	
HZ Herpes Zoster													■
Pneu-P-23 Pneumococcal Polysaccharide 23													■
Inf Influenza									* Every year in the fall				

◆ = A single vaccine dose given in a syringe and needle by intramuscular injection
 ■ = A single vaccine dose given in a syringe and needle by subcutaneous injection
 ▲ = A single vaccine dose given in an oral applicator by mouth
 ● = Provided through school-based immunization programs. Men-C-ACYW is a single dose; HB is a 2 dose series (see Table 6); HPV-4 is a 2 dose series (see Table 10). Each vaccine dose is given in a syringe and needle by intramuscular injection
 ^ = Preferably given at 4 years of age
 § = Given 10 years after the (4-6 year old) Tdap-IPV dose
 † = Given 10 years after the adolescent (14-16 year old) Tdap dose
 ‡ = Once a dose of Tdap is given in adulthood (24-26 years of age), adults should receive Td boosters every 10 years thereafter
 * = Children 6 months to 8 years of age who have not previously received a dose of influenza vaccine require 2 doses given ≥4 weeks apart. Children who have previously received ≥1 dose of influenza vaccine should receive 1 dose per season thereafter

Note: A different schedule and/or additional doses may be needed for high risk individuals (see Table 3) or if doses of a vaccine series are missed (see appropriate Tables 4-23)